

# Oregon Tuberculosis Control Program Indicators

## 1. Reporting and Treatment for TB cases

Outcome	Year					
	2000	2001	2002	2003	2004	2005
Completion of therapy <sup>†</sup>	81%	88%	87%	84%	90%	93%
HIV status reported	81%	76%	77%	77%	82%	93%
Totally directly observed therapy	55%	64%	68%	70%	61%	64%

## 2. Follow-up and Treatment for Contacts to Sputum-smear Positive Cases

Outcome	Year					
	2000	2001	2002	2003	2004	2005
Cases with no contacts	2%	11%	7%	6%	7%	6%
Contacts evaluated for TB	92%	84%	81%	88%	75%	71%
Active TB identified	2%	2%	1%	1%	0%	1%
NEW Latent TB identified <sup>‡</sup>	43%	39%	37%	32%	23%	19%
LTBI treatment started <sup>‡</sup>	71%	93%	92%	87%	80%	76%
LTBI treatment completed <sup>‡</sup>	Not Avail.	85%	73%	79%	58%	69%

Updated 12/2007

<sup>†</sup>Excludes: Patients with resistance to rifampin, patients who died during treatment, patients with meningeal TB, Children under age of 15 (at report) with disseminated TB (defined as military and or positive blood culture)

<sup>‡</sup> Excludes contacts with prior positive TST and/or prior history of treatment for TB infection or disease